

Content available at: <https://www.ipinnovative.com/open-access-journals>

IP Indian Journal of Clinical and Experimental Dermatology

Journal homepage: [www.ijced.org/](http://www.ijced.org/)

## Original Research Article

# The clinical type and etiological agents of superficial dermatophytosis: A cross sectional study

Shivam<sup>1</sup>, Shitij Goel<sup>1</sup>, Sushmita Agrahari<sup>1,\*</sup>, Gopi Krishna Maddali<sup>2</sup><sup>1</sup>Dept. of Dermatology, SMSR, Sharda University, Greater Noida, Uttar Pradesh, India<sup>2</sup>Heritage Institute of Medical Sciences, Varanasi, Uttar Pradesh, India

## ARTICLE INFO

## Article history:

Received 16-08-2021

Accepted 09-10-2021

Available online 11-12-2021

## Keywords:

Dermatophytic infections presenting with change in trend of etiological agent and becoming treatment unresponsive and recurrent.

## ABSTRACT

**Introduction:** Dermatophytosis are fungal infections caused by three genera of fungi that have the unique ability to invade and multiply within keratinized tissue (hair, skin, and nails). Although dermatomycoses are globally distributed, the endemic and most prevalent species of dermatophytosis differ strikingly from one geographic locality to another. Changing trend has been noticed in last few years with dermatophytic infections presenting as chronic, treatment unresponsive and recurrent. Also various microscopic and fungal culture studies have shown shift in identification of causative fungal species in recent years. Numerous studies have been done on the occurrence of dermatophytes in various parts of our country illustrating the range and changing pattern of fungal infection as well as causative fungal species.

**Materials and Methods:** Total number of 150 patients attending outpatient department of our hospital who were clinically diagnosed as having superficial dermatophytosis were enrolled into the study. Patients were carefully screened as per inclusion and exclusion criteria and then enrolled in the study. Samples were taken from all the patients and examined for KOH direct microscopy and sent for fungal culture on Sabouraud's Dextrose Agar as well as on Dermophyte Test Medium. Results were then analyzed using standard statistical methods.

**Results:** Out of total 150 patients, 101 were males and 49 were females. Most common age group was 21-30 years (37.3%). 58 patients (38.7%) showed positivity to KOH microscopy as well as fungal culture. Additionally 25 more samples demonstrated positivity to KOH microscopy (total 83 patients) but negativity to culture, while 9 patient samples were positive to culture but negative to direct microscopy. Predominant fungal species isolated on culture was *Trichophyton mentagrophytes* (50.7%) while next common species isolated was *T. tonsurans* (29.9%). No significant association was found between dermatophyte isolate on culture and clinical type. *Trichophyton mentagrophytes* and *Trichophyton tonsurans* were the most common species isolated among subjects with *Tinea faciei*, *Tinea cruris* and *Tinea corporis*.

**Conclusion:** The study showed a male preponderance and *T. corporis* was the commonest clinical type found. Majority of patients were in the 3rd decade and came within a duration of 1 month to 6 months of getting an infection. In patients diagnosed with *tinea corporis*, *tinea cruris*, *tinea pedis* and *tinea manuum*, *T. mentagrophytes* was the most predominant species isolated.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: [reprint@ipinnovative.com](mailto:reprint@ipinnovative.com)

## 1. Introduction

India is a big subcontinent with various weather as well as the topographic conditions. The hot and moist climate favours the acquisition and maintenance of fungal infections.<sup>1</sup> This in addition to congestion, inadequate

\* Corresponding author.

E-mail address: [liza.agrahari@gmail.com](mailto:liza.agrahari@gmail.com) (S. Agrahari).

socioeconomic conditions and poor hygiene enhances the chances of getting the fungal infection. Although dermatomycoses are globally distributed, the endemic and most prevalent species of dermatophytosis differ strikingly from one geographic locality to another.<sup>2</sup>

Numerous studies have been done on the occurrence of dermatophytes in various parts of our nation illustrating the range and changing pattern of fungal infection.<sup>1-5</sup> Superficial fungal infections are the most common fungal infections. According to World Health Organization (WHO), prevalence rate of superficial mycotic infection in children in developing countries has been found to be 20-25%.<sup>6</sup>

Dermatophytosis are fungal infections caused by three genera of fungi that have the unique ability to invade and multiply within keratinized tissue (hair, skin, and nails). These fungi, collectively called “dermatophytes”, are alike in their physiology, morphology, and pathogenicity. The three genera are *Microsporum*, *Trichophyton*, and *Epidermophyton*; species within these genera that do not invade keratinized tissue in animals or humans are not considered as dermatophytes. The identification of the fungal species is epidemiologically very crucial since the source of infection can be traced to prevent its transmission.<sup>7</sup> Direct microscopic examination is conducted making use of clearing solutions (KOH or Amman’s chloral lactophenol), while sensitivity may be enriched by utilizing stains/fluorochromes such as Congo red. Histological assessment does not allow accurate assessment of the fungal species.<sup>8</sup>

Culture is actually consequently required and specific culture media might be actually utilized to eliminate the growth of quickly growing contaminated moulds which might hinder the acknowledgment of dermatophytes. Several studies have been conducted in different parts of the country to study the changing trend in the causative organism of superficial dermatophytosis. We conducted this cross-sectional study in our patients to determine the clinical pattern of dermatophytosis prevalent in our center, to isolate and speciate the dermatophytes and to identify the change in trend in the causative organism of dermatophytosis.

## 2. Materials and Methods

After permission for institutional ethical committee a total of 150 consenting patients of superficial dermatophytosis presenting to the out patient department were taken in the study. Exclusion criteria was patients already on antifungals, patients on any oral immunosuppressant drug, patients with poorly controlled diabetes, immuno-compromised state like HIV, diabetes, previously treated patients, onychomycosis, candidal intertrigo, pityriasis versicolor, pregnant and lactating mothers. Complete history taking and detailed clinical examination was done in all the patients. Patients were explained everything about the study and procedure.

All the necessary laboratory investigations were performed in the study as per the study protocol. Specimen was collected from the affected area after it was cleaned with 70% ethyl alcohol and skin scales and crusts were collected in clean black paper. The specimen was further processed for KOH mount and culture on SDA and DTM.

For direct microscopy, the specimen was placed in a drop of 10% or 40% KOH on a microscopic slide, gentle heat was applied by passing the slide over a Bunsen flame for 3-4 times, the wet mount were covered with cover slip and left for 10-15 minutes before examination under microscope and the slides were examined under low power (10X) and high power (40X) magnification for demonstration of fungal elements. For culture on SDA and DTM, the specimen was inoculated on to two sets of test tubes, one containing Sabouraud’s dextrose agar with 0.05% chloramphenicol and 0.5% cycloheximide and the other on dermatophyte test medium. The specimen from skin were inoculated directly onto DTM and incubated at room temperature with the cap of the culture tube loose. Dermatophytes change the medium from yellow to red within 14 days.

Different deramtophytic species were identified as per their microscopic characteristics for example *Microsporum canis* spores are large, spindle-shaped, and thick-walled with six or more internal cells and often have a terminal knob. *Trichophyton mentagrophytes* was identified by long cigar-shaped macroconidia with thin walls. *Trichophyton rubrum* grows slowly in culture with sparse production of teardrop or peg-shaped microconidia laterally on fertile hyphae. Macroconidia, when present, are smooth-walled and narrowly club-shaped, although most isolates lack macroconidia.

For Microscopic examination of culture, the tease mount from the colony were prepared in lactophenol cotton blue and observed under low and high power objective of microscope, for the presence of hyphae, macroconidia, microconidia and other accessory structures of vegetative hyphae and the characters of each were noted.

### 2.1. Statistical analysis

The data was entered into the Microsoft excel and the statistical analysis was performed by statistical software SPSS version 25.0. The Quantitative (Numerical variables) were present in the form of mean and SD and the Qualitative (Categorical variables) were present in the form of frequency and percentage.

The chi-square test was applied for comparing the frequency between different groups. The p-value was considered to be significant when less than 0.05.

## 3. Results

Among the study population of 150, 101 were males and 49 females. The observed duration of illness was less than 1

month among 32 patients (21.3%), 1 month-6 months was reported among 62 (41.3%) and more than 6 months among 56 (37.3%) subjects. The clinical type of fungus species reported were Tinea faciei among 28 (18.7%), Tinea cruris among 84 (56.0%), Tinea corporis among 108 (72.0%), Tinea pedis among 7 (4.7%) and Tinea Manuum among 8 (5.3%) patients. There were 83 (55.3%) patient specimen which were KOH Positive, 67 (44.7%) Culture Positive and 58 (38.7%) both KOH and culture Positive. No significant association was found between Dermatophyte isolate on culture and clinical type. Trichophyton mentagrophytes and Trichophyton tonsurans was the most common species isolated among subjects with Tinea faciei, Tinea cruris and Tinea corporis.

**Table 1:** Age wise distribution of study population.

Age groups	Frequency	Percent
11-20 years	27	18.0%
21-30 years	56	37.3%
31-40 years	34	22.7%
41-50 years	19	12.7%
51-60 years	10	6.7%
61-70 years	4	2.7%

**Table 2:** Distribution of patients based on duration of illness.

Duration of illness	Frequency
< 1 month	32
1 month-6 months	62
> 6 months	56
Total	150


**Table 3:** Clinical type oftinea and their frequency in study population.

Clinical type	Frequency	Percent
Tinea faciei	28	18.7%
Tinea cruris	84	56.0%
Tinea corporis	108	72.0%
Tinea pedis	7	4.7%
Tinea manuum	8	5.3%

**Table 4:** Frequency of patients showing KOH positivity, Culture positivity or both.


	Frequency	Percent
KOH Positive	83	55.3%
Culture Positive	67	44.7%
Both KOH and culture Positive	58	38.7%

There were 83 (55.3%) KOH Positive, 67 (44.7%) Culture Positive and 58 (38.7%) Both KOH and culture Positive.




**Fig. 1:** Showing concordance and discordance between KOH and Fungal culture positivity

**Dermatophyte isolate on culture**



**Fig. 2:** Dermatophyte isolate on culture showed maximum result of Trichophyton mentagrophytes (50.7 %) followed by Trichophyton tonsurans (29.9%), Trichophyton rubrum (9.0%), epidermophyton flocculosum (7.5 %), Microsporum canis (3.0%).



**Fig. 3:** Comparison of distribution of culture positive and negative cases between different clinical types of fungus species.


**Table 5:** The distribution of culture positive and negative cases was compared between different clinical types of fungus species using the chi-square test. The culture positive cases were more among Tinea Manuum.

	Culture		Total	Chi-square value	p-value
	Negative	Positive			
Tinea faciei	17 60.7%	11 39.3%	28 100.0%	0.403	0.525
Tinea cruris	47 56.0%	37 44.0%	84 100.0%	0.030	0.863
Tinea corporis	60 55.6%	48 44.4%	108 100.0%	0.008	0.930
Tinea pedis	5 71.4%	2 28.6%	7 100.0%	0.770	0.380
Tinea manuum	7 87.5%	1 12.5%	8 100.0%	3.538	0.060

**Table 6:** Dermatophyte isolate on culture of different clinical types of tinea.

Dermatophyte isolate on culture	Clinical type				
	Tinea faciei	Tinea cruris	Tinea corporis	Tinea pedis	Tinea manuum
Epidermophyton Flocculosum	0	1	5	0	0
Microsporum canis	0	1	2	0	0
Trichophyton mentagrophytes	9	22	20	1	0
Trichophyton rubrum	2	3	3	0	1
Trichophyton tonsurans	0	10	18	1	0
	0.127	0.535	0.067	0.944	0.257

Trichophyton mentagrophytes was the most common species isolated among subjects with Tinea faciei, Tinea cruris and Tinea corporis.

**Fig. 4:** The distribution of KOH positive and negative cases was compared between different clinical types of fungus species using the chi-square test. The KOH positive cases were more among Tinea Faciei, Tinea cruris and Tinea corporis species though the difference was non-significant.

#### 4. Discussion

The studies on Dermatophytoses in India have received much attention in recent years because of the increasing incidence of the difficult to treat mycotic infections worldwide. Despite the current upsurge of dermatophytosis in India, it probably does not get the attention it deserves as far as scientific research is concerned. Antifungal susceptibility studies should be performed as increased

and prolonged use of antifungal agents has led to the development of resistance.<sup>9</sup>

In our study, the maximum number of subjects were found in the age groups of 21-30 and 31-40 years (37.3% and 22.7% respectively). Similar findings have been noted in studies done by Basak et al,<sup>10</sup> the maximum cases of dermatophytoses were seen in the age group of 21-30 years (25.4%) followed by age group of 31-40 years (20.3%), Walke HR et al.,<sup>11</sup> Poluri et al,<sup>12</sup> the highest incidence of dermatophytosis was observed in the age group of 21-40 years, Yadav et al,<sup>13</sup> dermatophytosis was more common in the age group of 21-30 years (30%) followed by 31-40 years (23%), Mishra M<sup>14</sup> and Sen and Raul.<sup>15</sup> Lower incidence in females could be attributed to their reluctance to seek medical advice, especially in rural areas where there may be a lack of female doctors or specialists.

The duration of presenting illness varied from less than 1 month in 21.3%, less than 1 month-6 months was reported among 41.3% and more than 6 months among 37.3% subjects. Our observations were slightly different from the study by Gupta et al,<sup>16</sup> where most of the patients had a prolonged duration of illness i.e. 58.8% patient had illness for more than 6 months. This was similar to the findings by Ghuse et al,<sup>17</sup> where the duration of infection was 1–2 months in the majority (43.3%) of the patients and only 6.7% had the disease for 1 year.

Tinea corporis (72.0%) was the most common clinical type found in the present study followed by Tinea cruris (56%). The other clinical types reported were Tinea faciei (18.7%), Tinea pedis (4.7%) and Tinea manuum (5.3%). This was in accordance with the findings of Basak et al,<sup>10</sup> Tinea corporis (52.65%) was the most common clinical presentation followed by Tinea unguium (14.1%) and Tinea cruris (12%) and Kucheria et al,<sup>18</sup> where the most common clinical presentation was Tinea corporis (31%) followed by Tinea unguium (21%). Studies by Walke HR et al<sup>11</sup> and Nagaral et al,<sup>19</sup> showed Tinea corporis as the most common clinical type followed by Tinea cruris. Agarwal et al<sup>20</sup> also reported that the most common clinical types observed was T. cruris (40%) followed by T. corporis (34.3%). However, studies by Gupta CM et al,<sup>21</sup> and Ghosh RR et al,<sup>22</sup> showed Tinea unguium as the most common clinical presentation followed by Tinea corporis and Tinea capitis.

In the current study, mixed infection was reported in majority (74.4%) of the samples among both males and females. Agrawalla et al<sup>23</sup> and Grando et al<sup>24</sup> conducted studies on anuum-mycological study of dermatophytosis observed that more than 20% of patients had two or more clinical types. In the present study, 55.3% samples were positive by direct microscopy by KOH mount, 44.7% samples were culture positive and 38.7% samples were both KOH positive and culture positive. The direct microscopy and culture findings of present study are relatively in agreement with study done by Dhyaneswari GP et al,<sup>25</sup> (72.6% KOH positive) and Mahale RP et al,<sup>26</sup> (61.01% culture positive).

In the study by Yadav et al,<sup>26</sup> out of 66 clinically diagnosed cases of dermatophytosis, 52 cases (78.79%) were positive for fungi, either by KOH and/or culture. 35 cases (53.03%) were positive by both KOH and culture, 13 cases (19.70%) were positive by KOH and negative by culture, 4 cases (6.06%) were negative by KOH but culture positive, 14 cases (21.21%) were negative by both KOH and culture, which is comparable with other studies done by Singh S<sup>1</sup> and Nada H.<sup>26</sup> This variation could be due to non-viability of fungal elements in some cases, inadequacy in sampling due to very small lesions and non-reported partial treatment with antifungal agents. In the present study, KOH positivity was 55.3% and culture positivity was 44.7% which was similar to the study by Poluri et al,<sup>12</sup> KOH positivity was 58.18% and culture positivity was 56.36%. However, similarity was seen with the study by Basak et al,<sup>10</sup> 71.1% samples were positive by direct microscopy by KOH mount and 59.8% samples were culture positive.

The culture positivity was found to be more in KOH positive cases with 38.7% cases being both KOH and culture positive. This shows that direct microscopy by KOH mount is a good screening test in the laboratory diagnosis of dermatophytosis.

The authors also came across instances where no fungal elements were seen under direct microscopy but showed growth on culture. This might be due to presence of scanty fungal elements which were missed during direct microscopic examination or due to presence of fungal elements in inactive sporulating form, which could not be visualised under direct microscopy.<sup>27,28</sup>

In our study, most commonly isolated dermatophyte species was Trichophyton mentagrophytes (50.7%) followed by Trichophyton tonsurans (29.9%), Trichophyton rubrum (9.0%), Epidermophyton Flocculosum (7.5%) and Microsporum canis (3.0%).

Our results were in accordance with the findings by Basak et al,<sup>10</sup> Trichophyton mentagrophytes (57.5%) was the most commonly isolated dermatophyte, followed by Trichophyton rubrum (30.1%), Trichophyton tonsurans (8.1%) and Microsporum anuum (4.3%), Gadadavar S et al,<sup>13</sup> (Trichophyton mentagrophytes 81.8%, Trichophyton rubrum 11.36%), Bhatia VK et al,<sup>29</sup> (Trichophyton mentagrophytes 63.5% and Trichophyton rubrum 35.1%) and Sahai et al.,<sup>30</sup> T. mentagrophytes was found to be the most common species.

This was contrasting to the research by Yadav et al,<sup>25</sup> T. rubrum (38.46%) was the commonest aetiological agent in majority of clinical types followed by T. mentagrophytes (33.33%), T. tonsurans (17.95%) and fungi other than Dermatophytes (10.26%) and Vineetha et al,<sup>14</sup> with T. rubrum was the most common species isolated in the first episode and in chronic dermatophytosis followed by T. mentagrophytes. Rare species, such as T. tonsurans and single cases of T. schoenleinii, E. floccosum and M. audouini, were isolated from chronic cases. Jain et al.,<sup>31</sup> Agarwal et al.<sup>20</sup> and Bindu et al.<sup>4</sup> also reported dissimilar findings to our study with T. rubrum being the most common species.

## 5. Conclusion

The study showed a male preponderance and T. corporis was commonest clinical type found. Majority of patients were in 3rd decade and came within a duration of 1 month to 6 months of getting infection. In patients diagnosed with tinea corporis, tinea cruris, tinea pedis and tinea manuum, T. mentagrophyte was the most predominant species isolated. Further research needs to be done to confirm the results.

Studies for antifungal susceptibility should be increased as dermatophytosis does not get the attention that it deserves even after the current changing trends of causative organisms and the resistance to various antifungals that it has developed.

## 6. Conflicts of Interest

The authors declare that there are no conflicts of interest regarding the publication of this paper.

## 7. Source of Funding

None.

## References

- Singh S, Beena PM. Profile of Dermatophytes infection in Baroda. *Indian J Dermatol Venereol Leprol.* 2003;69(4):281–3.
- Das K, Basak S, Ray S. A Study on Superficial fungal infection from west Bengal: A brief report. *J Life Sci.* 2009;1(1):51–5. doi:10.1080/09751270.2009.11885134.
- Grove S, Roy P. Clinico-mycological Profile of Superficial Mycosis in a Hospital in North-East India. *Med J Armed Forces India.* 2003;59(2):114–6. doi:10.1016/S0377-1237(03)80053-9.
- Bindu V, Pavithran K. Clinico-mycological study of dermatophytosis in Calicut. *Indian J Dermatol Venereol Leprol.* 2002;68(5):259–61.
- Gopi A, Harindranath D, Kaushik AR. Mycological profile of dermatophytes isolated from clinical samples in KIMS hospital Bangalore. *J Evol Med Dent Sci.* 2015;4(2):835–42.
- WHO, 2005. Epidemiology and management of common skin diseases in children in developing countries. World Health Organization, Geneva. WHO/FCH/CAH/05.12; 2005.
- Gupta AK, Summerbell RC. Increased incidence of Trichophyton tonsurans tinea capitis in Ontario, Canada between 1985 and 1996. *Med Mycol.* 1998;36:55–60.
- Shalaby MF, El-Din AN, El-Hamd MA. Isolation, identification, and in vitro antifungal susceptibility testing of dermatophytes from clinical samples at Sohag University Hospital in Egypt. *Electron Physician.* 2016;8(6):2557–67. doi:10.19082/2557.
- Jain S, Kabi S, Swain B. Current Trends of Dermatophytosis in Eastern Odisha. *J Lab Physicians.* 2020;12(1):10–4. doi:10.1055/s-0040-1713063.
- Basak P, Mallick B, Pattanaik S. Prevalence of dermatophytic infections including antifungal susceptibility pattern of dermatophytes in a tertiary care hospital. *Int J Res Med Sci.* 2019;7(3):699–705. doi:10.18203/2320-6012.ijrms20190461.
- Walke HR, Gaikwad AA, Palekar SS. Clinicomycological profile of dermatophytosis in patients attending dermatology OPD in tertiary care hospital, India. *Int J Curr Microbiol App Sci.* 2014;3(10):432–40.
- Poluri LV, Indugula JP, Kondapaneni SL. Clinicomycological Study of Dermatophytosis in South India. *J Lab Physicians.* 2015;7(2):84–9.
- Lyngdoh CJ, Lyngdoh WV, Choudhury B, Sangma KA, Bora I, Khyriem AB, et al. Clinico-mycological profile of dermatophytosis in Meghalaya. *Int J Med Public Health.* 2013;3(4):254–6.
- Mishra N, Rastogi MK, Gahalaut P, Yadav S, Srivastava N, Aggarwal A, et al. Clinicomycological study of dermatophytoses in children: Presenting at a tertiary care center. *Indian J Paediatr Dermatol.* 2018;19(4):326–30. doi:10.4103/ijpd.IJPD\_98\_17.
- Sen SS, Raul ES. Dermatophytosis in Assam. *Indian J Med Microbiol.* 2006;24(1):77–8. doi:10.4103/0255-0857.19907.
- Gupta AK, Mohan A, Singh SK, Pandey AK. Studying the clinic mycological pattern of the dermatophytic infection attending OPD in tertiary care hospital in eastern Uttar Pradesh and Bihar. *Int J Res Dermatol.* 2018;4(2):118–25. doi:10.18203/issn.2455-4529.
- Ghuse V, Someshwar S, Jerajani H. Patterns of culture positivity and antifungal sensitivity in dermatophytosis. *MGM J Med Sci.* 2019;6(3):105–12. doi:10.4103/mgmj.mgmj\_4\_20.
- Kucheria M, Gupta SK, Chinna DK, Gupta V, Hans D, Singh K, et al. Clinicomycological Profile of Dermatophytic Infections at a Tertiary Care Hospital in North India. *Int J Com Health and Med Res.* 2016;2(2):17–22.
- Nagaral GV, Goud GK, Sudha P. Prevalence of tinea corporis and tinea cruris in Chitradurga rural population. *Indian J Clin Exp Dermatol.* 2018;4(3):221–5.
- Agarwal U, Saran J, Agarwal P. Clinico-mycological study of dermatophytes in a tertiary care centre in northwest India. *Indian J Derm Venereol Leprol.* 2014;80(2):194. doi:10.4103/0378-6323.129434.
- Gupta CM, Tripathi K, Tiwari S, Rathore Y, Nema S, Dhanvijay AG, et al. Current trends of clinicomycological profile of dermatophytosis in Central India. *J Dent Med Sci.* 2014;13(10):23–6.
- Ghosh RR, Ray R, Ghosh TK, Ghosh AP. Clinicomycological profile of dermatophytoses in a tertiary care centre hospital in West Bengal-An Indian Scenario. *Int J Curr Microbiol App Sci.* 2014;3(9):655–6.
- Agarwalla A, Jacob M, Sethi M, Parija SC, Singh NP. A clinico-mycological study of dermatophytoses in Nepal. *J Dermatol.* 2001;28(1):16–21.
- Grando SA, Hostager BS, Herron MJ, Dahl MV, Nelson RD. Binding of Trichophyton rubrum Mannan to Human Monocytes In Vitro. *J Invest Dermatol.* 1992;98(6):876–80. doi:10.1111/1523-1747.ep12457923.
- Dhyaneswari GP, Muley VA, Bhore AV. Clinicomycological profile of dermatophytosis in a tertiary care hospital in Western India. *SAS J Med.* 2015;1(4):160–5.
- Mahale RP, Rao MR, Tejashree A, Deepashree R, Kulkarni M. Clinicomycological profile of dermatophytosis in a teaching hospital. *Int J Pharmaceut Sci Invent.* 2014;3(8):43–49.
- History of antifungals. Department of Dermatology. *J Am Acad.* 1990;4(2):776–784.
- Roopa C, Biradar S. Incidence and Identification of Dermatophytes in a Tertiary Care Hospital in North Karnataka, India. *Int J Curr Microbiol App Sci.* 2015;4(9):986–90.
- Gadadavar S, Shilpa HS, Patil CS, Vinay PS, Shettar N. Clinico-Mycological study of dermatophytoses at a tertiary care hospital in Belagavi. *Int J Curr Microbiol App Sci.* 2018;7(5):1872–80.
- Bhatia VK, Sharma PC. Epidemiological studies on dermatophytosis in human patients in Himachal Pradesh, India. *Springerplus.* 2014;3(1):134. doi:10.1186/2193-1801-3-134.
- Jain N, Sharma M, Saxena V. Clinico-mycological profile of dermatophytosis in Jaipur, Rajasthan. *Indian J Dermatol Venereol Leprol.* 2008;74(3):274–5. doi:10.4103/0378-6323.41388.

## Author biography

**Shivam**, Resident

**Shitij Goel**, Professor

**Sushmita Agrahari**, Resident

**Gopi Krishna Maddali**, Professor and Head

**Cite this article:** Shivam, Goel S, Agrahari S, Maddali GK. The clinical type and etiological agents of superficial dermatophytosis: A cross sectional study. *IP Indian J Clin Exp Dermatol* 2021;7(4):331-336.