



## Original Research Article

# A cross sectional study optimizing platelet rich plasma preparation using various centrifugation speeds and its effects on the platelet yield

Zeal Jayeshbhai Rachchha<sup>1</sup>, Vaishali Sanket Makwana<sup>1\*</sup>,  
Sandip Labhubhai Lathiya<sup>1</sup>, Nehalben Manishbhai Pandya<sup>1</sup>

<sup>1</sup>Dept. of Dermatology, C.U. Shah Medical College and Hospital, Surendranagar, Gujarat, India



## ARTICLE INFO

## Article history:

Received 15-02-2024

Accepted 20-05-2024

Available online 04-09-2024

## Keywords:

Plateletrich plasma

PRP

Centrifugation speed

Platelet count

## ABSTRACT

**Background:** Platelet-rich plasma (PRP) is an autologous preparation of platelets in concentrated plasma (with usually >1,000,000 platelets/ $\mu$ L or 2–7 times the concentration in whole blood), prepared by a process known as differential centrifugation. The therapeutic efficacy of a PRP preparation is largely determined by its platelet concentration. Although PRP has been used for many years, there is no standard preparation protocol. This study was conducted to compare different protocols to determine which spin variation produces the highest platelet yield.

**Materials and Methods:** A cross sectional study was conducted at a tertiary care hospital with 35 patients. We included the data of the patients that were enrolled in PRP therapy for various dermatological indications. We analysed the data for PRP prepared using different centrifugation protocols and then compared the platelet count of PRP with the baseline platelet count of whole blood to obtain the platelet yield. The platelet yields for different protocols were compared using statistical analysis (IBM SPSS software).

**Results:** After comparing the platelet yield of different protocols, we observed that centrifugation at 1300rpm/246 g for 20 minutes followed by 1600 rpm/373 g for 20 minutes achieved the highest platelet yield-4.6468 times as compared to the mean platelet count in the whole blood and the difference was statistically significant.

**Conclusion:** Multiple parameters influence the platelet concentration obtained in PRP. Therefore, the practitioner must consider the method of PRP preparation that gives the optimum platelet yield to deliver the best therapeutic results to the patients.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: [reprint@ipinnovative.com](mailto:reprint@ipinnovative.com)

## 1. Introduction

Platelet-rich plasma (PRP) is an autologous preparation of platelets in concentrated plasma (with usually >1,000,000 platelets/ $\mu$ L or 2–7 times the concentration in whole blood).<sup>1</sup> PRP contains various growth factors in the alpha granules of platelets, such as platelet-derived growth factor, vascular endothelial growth factor, epithelial growth factor, fibroblast growth factor, hepatic growth factor, transforming

growth factor, etc. These growth factors play a role in tissue repair, regeneration, angiogenesis, connective tissue remodelling and cell cycle regulation.<sup>2</sup>

The method known as differential centrifugation is used for PRP preparation. Apart from dermatology, PRP is used in various medical specialities like orthopaedics, maxillofacial surgery, regenerative medicine, gynaecology, dentistry and sports medicine.<sup>3,4</sup> In dermatology, PRP is used in alopecia (angiogenic role), skin rejuvenation (increases dermal elasticity), acute and chronic ulcers (enhanced re-epithelisation by growth factors), acne

\* Corresponding author.

E-mail address: [123vaishali123@gmail.com](mailto:123vaishali123@gmail.com) (V. S. Makwana).

scars and post-surgical scars(collagen synthesis), lipodermatosclerosis, etc.<sup>5</sup>

Although PRP has been used for many years, no standard preparation protocol exists. Therefore, attaining a count of >1,000,000platelets/ $\mu$ L without using expensive PRP kits in a clinic or even a tertiary care setting remains a hurdle. Hence, this study aims to analyze the platelet count of PRP prepared using various centrifugation parameters to achieve the ideal PRP.

## 2. Materials and Methods

We conducted this cross sectional study at the dermatology outpatient department of a tertiary care hospital from data of the period of 10 months from March 2022 to December 2022. The study was approved by the Institutional Ethics Committee of the Hospital. A data of 35 apparently healthy individuals who were enrolled for PRP therapy for various dermatological indications like androgenic alopecia, acne scars, melasma, etc. was collected.

### 2.1. Inclusion criteria

Patients of all age groups and both genders enrolled for PRP therapy for various dermatological indications e.g. androgenic alopecia, acne scar and melasma.

### 2.2. Exclusion criteria

None.

### 2.3. Statistical analysis

The data was analyzed using statistical software - SPSS version 26 (IBM Corp. released 2019. IBM SPSS Statistics for Windows, Version 26.0 Armonk, NY: IBM Corp.). The platelet yields derived from different centrifugation speeds were compared using non-parametric tests (Kolmogorov Smirnov test & Shapiro Wilk test).

Charts and tables were prepared using Microsoft Word@ 2010, Microsoft Office Professional Plus, and Microsoft Corp.

Sample size: Total- 35




Chart 1: Study methodology flowchart

### 2.4. Procedure of PRP preparation

The whole procedure of PRP preparation was carried out in an air-conditioned environment at 24 degrees Celsius.

The rotor's size and radius (R) vary with different centrifuge machines. Therefore, relative centrifugal force (g) values are converted to rotation per minute (rpm). The conversion factor from 'g' to rpm is as follows<sup>4</sup>:

$$g = (1.118 \times 10^{-5}) R (\text{rpm})^2$$

Peripheral blood (20 cc) was obtained from each patient, out of which 3ml was added to an EDTA (Ethylenediaminetetraacetic acid) tube for baseline platelet count, and the rest of it was divided into 2 BD (Becton Dickinson) vacutainer whole blood tube (Acid Citrate Dextrose-A, Yellow, 8.5 ml) for PRP preparation. PRP was prepared in a Remi Medico Plus centrifuge machine (Figure 1)(Radius of centrifuge-13 cm) using a two-step centrifugation method.



Figure 1: Commercially available table top remi medico plus centrifuge machine


The first spin is performed according to the centrifugation speed and time allotted for that subject. After the first spin, the whole blood separates into three layers: an upper layer containing platelet-poor plasma (PPP), an intermediate layer called buffy coat rich in WBCs and platelets and a bottom layer containing RBCs as shown in Figure 3. The upper layer, along with the buffy coat, is transferred into a sterile plain vacutainer and placed in the centrifuge machine for the second spin.

The second spin is performed with predetermined centrifugation speed and time. The resultant plasma contains platelet-poor plasma (PPP) in the upper 3/4<sup>th</sup> volume and platelet-rich plasma (PRP) in the lower 1/4<sup>th</sup> volume, as shown in Figure 4.


The upper 3/4<sup>th</sup> portion (PPP) is withdrawn with the syringe and discarded, while the lower 1/4<sup>th</sup> portion (PRP) is resuspended to form a homogenized PRP, as shown in Figure 5.

**Table 1:** Protocols of PRP preparation

Spin Variations	1 <sup>st</sup> centrifugation speed (Rotation per minute-rpm)	1 <sup>st</sup> centrifugation speed (Relative centrifugal force- g force)	Time (minutes)	2 <sup>nd</sup> centrifugation speed (Rotation per minute-rpm)	2 <sup>nd</sup> centrifugation speed (Relative centrifugal force- g force)	Time (minutes)
1	1020	151 g	20	1780	461 g	10
2	1020	151 g	10	1020	151 g	10
3	1300	246 g	20	1020	151 g	20
4	1500	328 g	20	1120	183 g	20
5	1500	328 g	20	1150	193 g	20
6	1500	328 g	20	1200	210 g	20
7	1300	246 g	20	1600	373 g	20




**Figure 2:** Whole blood collected in 2 ACDA (Acid Citrate Dextrose-A) vacutainer




**Figure 3:** After 1<sup>st</sup> spin, supernatant plasma, buffy coat and RBC layer are formed

The seven protocols selected were inspired by the previous studies with few modifications, which were conducted to determine optimum centrifugation speeds for PRP preparation.<sup>6-11</sup> The data from seven groups containing an equal number of samples (5 samples in each group) were collected in this study. Various centrifugation protocols used in our study are given in Table 1.



**Figure 4:** After 2<sup>nd</sup> spin, upper 3/4<sup>th</sup> PPP is formed and lower 1/4<sup>th</sup> PRP is formed




**Figure 5:** Upper 3/4<sup>th</sup> PPP is discarded, and lower 1/4<sup>th</sup> PRP containing platelet pellet and RBC at the bottom is resuspended to form homogenized PRP

In each of the study groups, the mean platelet count of the whole blood and PRP are calculated, and the platelet yield is obtained.

$$\text{Platelet yield} = \frac{\text{mean platelet count in Platelet-rich plasma}}{\text{mean baseline platelet count in the whole blood}}$$

### 3. Results

Table 2 depicts the baseline platelet count, PRP platelet count and platelet yield, as well as the averages for all seven spin variations used in our study. The data consisted of 35 samples from 35 individuals, comprising 40% males (n=14) and 60% females (n=21), as shown in Figure 6. They were in the age group of 18–40 years. Androgenic alopecia was the most common indication, followed by acne scar.



**Figure 6:** Gender distribution among the study participants

In our study, the highest mean platelet count was achieved by applying spin variation-7 (1300 rpm/246 g for 20 minutes followed by 1600 rpm/373 g for 20 minutes), which was 13,31,400/cumm. The second highest mean platelet count was achieved by spin variation - 1 (1020 rpm/151 g for 20 minutes followed by 1780 rpm/461 g for 10 minutes), which was 9,91,200/cumm as shown in Figure 7. Therefore, the highest platelet yield is achieved by spin variation-7, which is 4.6468, followed by spin variation- 1, which is 3.2068, as shown in Figure 8.

The platelet yields derived from different centrifugation speeds were compared with each other using non-parametric tests (Kolmogorov Smirnov test & Shapiro Wilk test). Since the sample size is small (< 40) so, data is checked for Shapiro Wilk significance. It is >0.05, so data follows a normal distribution, as shown in Table 3.

One-way Analysis of Variance (ANOVA) was used for analysis as data followed a normal distribution. It is followed by Tukey’s Honestly Significant Difference (HSD) post hoc test for multiple individual comparisons.

**Table 2:** Results obtained in various spin variations

Spin variation	Rotation		Baseline platelet count (/cumm)	PRP platelet count (/cumm)	Platelet yield (× times)	Average platelet yield	
	1 <sup>st</sup> rotation	2 <sup>nd</sup> rotation					
1	1020 rpm for 20 minutes	1780 rpm for 10 minutes	273000	1082000	3.963	3.2068	
			246000	763000	3.102		
			262000	721000	2.752		
			403000	1576000	3.911		
			353000	814000	2.306		
			<b>Average</b>	<b>307400</b>	<b>991200</b>		<b>3.2068</b>
	321000	741000	2.308				
2	1020 rpm for 10 minutes	1020 rpm for 10 minutes	378000	672000	1.777	2.4306	
			321000	673000	2.097		
			211000	548000	2.597		
			262000	884000	3.374		
			<b>Average</b>	<b>298600</b>	<b>703600</b>		<b>2.4306</b>
			254000	917000	3.61		
	253000	772000	3.051				
3	1300 rpm for 20 minutes	1020 rpm for 20 minutes	162000	564000	3.481	3.1488	
			287000	790000	2.753		
			378000	1077000	2.849		
			<b>Average</b>	<b>266800</b>	<b>824000</b>		<b>3.1488</b>
			373000	858000	2.3		
			395000	937000	2.372		
	246000	571000	2.321				
4	1500 rpm for 20 minutes	1120 rpm for 20 minutes	211000	566000	2.682	2.6412	
			273000	964000	3.531		
			<b>Average</b>	<b>299600</b>	<b>779200</b>		<b>2.6412</b>
			272000	765000	2.812		
			296000	876000	2.959		
			222000	625000	2.815		
	374000	1052000	2.813				
5	1500 rpm for 20 minutes	1150 rpm for 20 minutes	230000	602000	2.617	2.8032	
			<b>Average</b>	<b>278800</b>	<b>784000</b>		<b>2.8032</b>
			296000	1084000	3.662		
			287000	735000	2.561		
			291000	858000	2.948		
			269000	777000	2.888		
	373000	1113000	2.984				
6	1500 rpm for 20 minutes	1200 rpm for 20 minutes	353000	1436000	4.068	3.0086	
			<b>Average</b>	<b>303200</b>	<b>913400</b>		<b>3.0086</b>
			162000	879000	5.426		
			324000	1755000	5.417		
			373000	1496000	4.011		
			253000	1091000	4.312		
	<b>Average</b>	<b>293000</b>	<b>1331400</b>	<b>4.6468</b>			
7	1300 rpm for 20 minutes	1600 rpm for 20 minutes	162000	879000	5.426	4.6468	
			324000	1755000	5.417		
			373000	1496000	4.011		
			253000	1091000	4.312		
			<b>Average</b>	<b>293000</b>	<b>1331400</b>		<b>4.6468</b>
			162000	879000	5.426		
	324000	1755000	5.417				

The test statistic is  $F_{(6, 28)} = 9.219$  ( $p = .000$ ). Using an  $\alpha$  of 0.05, we get a critical value of  $F_{0.05;(6,28)} = 2.445$ . Since the test statistic is much larger than the critical value, we reject the null hypothesis and conclude that a statistically significant difference exists. There was a statistically significant difference between spin variation - 7 when compared to spin variations - 1 to 6 as determined by one-way ANOVA, as shown in Table 5.

A Tukey post hoc test revealed that the simultaneous pairwise comparisons between spin variations - 1 to 6 are

not significant ( $p \geq 0.05$ ), whereas the comparison between spin variation - 7 with respect to other spin variations is significant ( $p < 0.05$ ), as shown in Table 6. Thus, platelet yield obtained was statistically significantly higher with spin variation-7 after comparing spin variations - 1 to 6 with each other as well as with spin variation-7 ( $4.6468 \pm 0.71619$ ) ( $p = .000$ ). There was no statistically significant difference found during multiple comparisons between the spin variations - 1 to 6.



**Table 3:** Tests of normality of data

Methods	Tests of Normality						
	Kolmogorov-Smirnov <sup>a</sup>				Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.	
platelet yield	1	.235	5	.200*	.910	5	.465
	2	.192	5	.200*	.950	5	.736
	3	.209	5	.200*	.900	5	.412
	4	.297	5	.170	.755	5	.099
	5	.329	5	.082	.878	5	.298
	6	.324	5	.092	.882	5	.319
	7	.280	5	.200*	.781	5	.057

\*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

(df: degree of freedom; Sig: Significance)

**Table 4:** Descriptive data for all seven different centrifugation speeds

Methods	N	Mean	Standard Deviation	Descriptives Platelet Yield				Minimum	Maximum
				Standard Error	95% Confidence Interval for Mean				
					Lower Bound	Upper Bound			
1	5	3.2068	.72405	.32381	2.3078	4.1058	2.31	3.96	
2	5	2.4306	.60647	.27122	1.6776	3.1836	1.78	3.37	
3	5	3.1488	.38051	.17017	2.6763	3.6213	2.75	3.61	
4	5	2.6412	.52077	.23290	1.9946	3.2878	2.30	3.53	
5	5	2.8032	.12171	.05443	2.6521	2.9543	2.62	2.96	
6	5	3.0086	.40190	.17974	2.5096	3.5076	2.56	3.66	
7	5	4.6468	.71619	.32029	3.7575	5.5361	4.01	5.43	
Total	35	3.1266	.83616	.14134	2.8393	3.4138	1.78	5.43	

**Table 5:** ANOVA test

	ANOVA Platelet Yield				
	Sum of Squares	df	Mean Square	F	Sig.
Between Groups	15.782	6	2.630	9.219	.000
Within Groups	7.989	28	.285		
Total	23.772	34			

(ANOVA – Analysis of Variance, df – degree of freedom, F – Variance)

#### 4. Discussion

PRP was first described in 1997 by Whitman et al. as a derivative of the fibrin glue.<sup>12</sup> The therapeutic efficacy of a PRP preparation is largely determined by its platelet concentration. Therefore, clinicians should be aware of the fact that different PRP preparation protocols can produce different platelet concentrations, which can translate into a difference in efficacy and clinical response.<sup>13</sup> As per the study conducted by Dashore S et al., the recommended parameters of centrifugation for PRP preparation are 100–300 g for 5–10 minutes for the first spin and 400–700 g for 10–17 minutes for the second spin.<sup>4</sup> Various factors like the type of anti-coagulant, centrifugal speeds, the amount and the


kind of growth factors existing in PRP, the number of platelets in the donor's blood and PRP are essential for determining the procedure's efficacy and clinical outcome. Therefore, they must be carefully considered before applying to clinical practice.<sup>10,14–16</sup> Haynesworth et al. demonstrated that the proliferation and differentiation of adult mesenchymal stem cells were directly related to platelet concentration. A concentration of approximately 4 to 5 times the baseline platelet count would be required to produce a sufficient cellular response.<sup>17</sup> Rughetti et al. found that there is a bell-curve relationship between the platelet count and its functional activity. Therefore, the optimal proliferation and differentiation of endothelial cells occurred at  $1.25 \times 10^6$  platelets/ml, and optimal angiogenesis occurred at  $1.5 \times 10^6$  platelets/ml. As the platelet count

**Table 6:** Tukey’s HSD post-hoc test


Multiple Comparisons							
Dependent Variable: platelet yield							
Tukey’s HSD post hoc test							
(I) Methods	(J) Methods	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval		
					Lower Bound	Upper Bound	
1	2	.77620	.33783	.280	-.2955	1.8479	
	3	.05800	.33783	1.000	-1.0137	1.1297	
	4	.56560	.33783	.638	-.5061	1.6373	
	5	.40360	.33783	.890	-.6681	1.4753	
	6	.19820	.33783	.997	-.8735	1.2699	
	7	-1.44000*	.33783	.003	-2.5117	-.3683	
	1	2	-.77620	.33783	.280	-1.8479	.2955
2	3	-.71820	.33783	.366	-1.7899	.3535	
	4	-.21060	.33783	.995	-1.2823	.8611	
	5	-.37260	.33783	.922	-1.4443	.6991	
	6	-.57800	.33783	.615	-1.6497	.4937	
	7	-2.21620*	.33783	.000	-3.2879	-1.1445	
	1	2	-.05800	.33783	1.000	-1.1297	1.0137
	1	3	.71820	.33783	.366	-.3535	1.7899
3	4	.50760	.33783	.741	-.5641	1.5793	
	5	.34560	.33783	.944	-.7261	1.4173	
	6	.14020	.33783	1.000	-.9315	1.2119	
	7	-1.49800*	.33783	.002	-2.5697	-.4263	
	1	2	-.56560	.33783	.638	-1.6373	.5061
	1	3	.21060	.33783	.995	-.8611	1.2823
	1	4	-.50760	.33783	.741	-1.5793	.5641
4	5	-.16200	.33783	.999	-1.2337	.9097	
	6	-.36740	.33783	.927	-1.4391	.7043	
	7	-2.00560*	.33783	.000	-3.0773	-.9339	
	1	2	-.40360	.33783	.890	-1.4753	.6681
	1	3	.37260	.33783	.922	-.6991	1.4443
	1	4	-.34560	.33783	.944	-1.4173	.7261
	1	5	.16200	.33783	.999	-.9097	1.2337
5	6	-.20540	.33783	.996	-1.2771	.8663	
	7	-1.84360*	.33783	.000	-2.9153	-.7719	
	1	2	-.19820	.33783	.997	-1.2699	.8735
	1	3	.57800	.33783	.615	-.4937	1.6497
	1	4	-.14020	.33783	1.000	-1.2119	.9315
	1	5	.36740	.33783	.927	-.7043	1.4391
	1	6	.20540	.33783	.996	-.8663	1.2771
6	7	-1.63820*	.33783	.001	-2.7099	-.5665	
	1	2	1.44000*	.33783	.003	.3683	2.5117
	1	3	2.21620*	.33783	.000	1.1445	3.2879
	1	4	1.49800*	.33783	.002	.4263	2.5697
	1	5	2.00560*	.33783	.000	.9339	3.0773
	1	6	1.84360*	.33783	.000	.7719	2.9153
	1	7	1.63820*	.33783	.001	.5665	2.7099

\*. The mean difference is significant at the 0.05 level.

(Tukey’s HSD post hoc test- Tukey’s Honestly Significant Difference post hoc test)



**Figure 7:** Graph showing mean baseline platelet count versus mean platelet count in PRP in each study group



**Figure 8:** Platelet yield in different spin variations used in our study

increased higher than  $1.5 \times 10^6$  platelets/ml, the proliferation of endothelial cells showed a decreasing trend.<sup>18</sup>

A classification of PRP preparation was proposed by Dohan Ehrenfest DM, Rasmusson L, Albrektsson T in 2009 as follows:<sup>19</sup>

1. Pure platelet-rich plasma (P-PRP) or leukocyte-poor platelet-rich plasma: It contains no WBC and low-density fibrin network after activation.
2. Leukocyte and platelet-rich plasma (L-PRP): It contains WBCs and a low-density fibrin network after activation. Most of the commercially available systems belong to this group.
3. Pure platelet-rich fibrin (P-PRF)/ leukocyte poor platelet-rich fibrin preparation: It contains no WBC and a high-density fibrin network. They exist only in strongly activated gel forms and cannot be injected like traditional fibrin glues.
4. Leukocyte and platelet-rich fibrin (L-PRF)/second-generation platelet-rich plasma: It contains WBC along with a high-density fibrin network.

The buffy coat inclusion methods give a higher platelet concentration in PRP, but the disadvantage of this method is higher contamination with RBCs and WBCs.<sup>3</sup> The role of WBCs in PRP is not clear. It is believed that WBCs in PRP provide protection from infections, increase the release of growth factors and help in angiogenesis.<sup>19</sup> Also, as mentioned by Oudelaar et al., the concentration of vascular endothelial growth factor is significantly higher in PRP produced by systems with higher concentrations of platelets and leucocytes than in P-PRP kits.<sup>20</sup> In PRP, RBCs may affect platelet function by altering pH and promoting inflammation. Also, RBCs can act as a source of reactive oxygen species. When used for skin rejuvenation, RBCs may also induce post-inflammatory hyperpigmentation due to hemosiderin deposition.<sup>21–23</sup> But RBC contamination is quite unavoidable in the PRP preparation, including the buffy coat layer, even if care is taken while separating the buffy coat from a layer from RBC volume. There is no consensus on whether or not the platelet must be previously activated before their application and which activator to use. In some studies, platelets are activated using thrombin or calcium, while in other studies; platelets are not activated.<sup>24</sup>

In today’s era, numerous centrifugation methods are available for PRP preparation. However, there is a need for more literature to determine the best method of PRP preparation. Some authors recommend the single spin method wherein the whole blood is centrifuged only once. Kahn RA et al. showed that centrifugation of 3731 g for 478 ml of whole blood for a period of 4 minutes was optimal to obtain the highest platelet concentration in the sample<sup>8</sup>. However, many studies state that a double spin method of PRP preparation yields a higher platelet count. A comparison of our research with various protocols for PRP preparation by different authors is mentioned in Table 7.

In our study, we have used the data of protocols with slightly modified spin parameters used in previous studies and studied the data of seven novel spin parameters. We have used the data of buffy coat inclusion method in our study. We compared the platelet yield instead of the mean platelet count obtained by each group so that we could avoid the bias created by the difference in the baseline platelet count of participants of each group. Here, the highest mean platelet yield is achieved by applying spin variation-7 (1300 rpm/246 g for 20 minutes followed by 1600 rpm/373 g for 20 minutes), which was 13,31,400/cumm (4.6468 times as compared to mean platelet count in the whole blood).

In our study, spin variation-7 produced the mean platelet count of PRP of more than 1,000,000/cumm, which, as described earlier, is required for producing optimal results by PRP. The spin variations 1, 2, 3, 4, 5 and 6 have produced a platelet count of less than 1,000,000/cumm, so they may not produce the desired clinical outcomes.

Araki et al. used varying accelerations ranging from 230-270g in the first spin and 2330g in the second spin. They



**Table 7:** Comparison of various protocols of PRP preparation

S.No.	Study	1 <sup>st</sup> Spin	2 <sup>nd</sup> spin	Platelet concentration in PRP (x10 <sup>6</sup> /cumm)	% of the increase in platelet counts (times)
1	Araki et al <sup>25</sup>	230 – 270 g x 10 minutes	2330 g x 10 minutes	1.89	7.4
2	Amable et al <sup>7</sup>	300 g x 5 minutes	700 g x 17 minutes	0.14 – 0.19	5.4 – 7.3
3	Tamimi et al <sup>26</sup>	160 g x 10 minutes	400 g x 10 minutes	0.632	3.52
4	Bausset et al <sup>27</sup>	130 g x 15 minutes	250 g x 15 minutes	-	3.96
5	Our study	246 g x 20 minutes	373 g x 20 minutes	1.47	4.6468

achieved the mean platelet count of  $1.89 \times 10^6$  /cumm and a platelet yield of 7.4 times as compared to the mean platelet count of PRP of  $1.33 \times 10^6$  /cumm and platelet yield of 4.6468 times in our study.<sup>25</sup> Here, the first spin speed was comparable to our study, but the second spin speed was very high compared to our study. Also, the time used for PRP preparation is less compared to our study. However, few authors believe that the longer time periods slightly increase the platelet recovery and decrease the WBC in the upper layer. Hence, time can be a control parameter when low levels of WBCs are required in the PRP sample.<sup>24</sup>

In our study, a higher spin speed is applied for the second spin than the first spin, which is in accordance with the other studies compared here.

To our knowledge, we found that there were many methods for preparing PRP, but no comparative study similar to the one shown in the present report has been described previously. The limitation of our study was a limited sample size; hence, further studies with larger sample sizes are required.

## 5. Conclusion

The science behind the preparation of PRP is still evolving. There is no single "ideal" method for the preparation of PRP. Multiple factors like temperature, cell counts of whole blood, centrifugation method parameters, and centrifuge machine parameters play a significant role in the quality of PRP prepared. In our study, the highest platelet count in PRP is achieved by spin variation-7 (1300 rpm/246 g for 20 minutes followed by 1600 rpm/373 g for 20 minutes), which was 13,31,400/cumm (4.6468 times as compared to the mean platelet count in the whole blood). It is necessary that dermatologist take into account various facets of PRP preparation to ensure that the best therapeutic result is delivered to their patients.

## 6. Conflict of Interest

None.

## 7. Source of Funding

None.

## Acknowledgements

None.

## References

- Marx RE. Platelet-rich plasma (PRP): what is PRP and what is not PRP? *Implant Dent.* 2001;10(4):225–8.
- Kaushik A, Kumaran MS. Platelet-Rich Plasma: The Journey so Far! *Indian. Dermatol Online J.* 2020;11(5):685–92.
- Muthuprabakaran K, Pai VV, Ahmad S, Shukla P. A cross-sectional analysis of the effects of various centrifugation speeds and inclusion of the buffy coat in platelet-rich plasma preparation. *Indian J Dermatol Venereol Leprol.* 2021;87(6):792–9.
- Dashore S, Chouhan K, Nanda S, Sharma A. Preparation of Platelet-Rich Plasma: National IADVL PRP Taskforce Recommendations. *Indian Dermatol Online J.* 2021;12(1):12–23.
- Mohan S, Mohan L, Sangal R, Singh N. Platelet rich plasma in dermatology and cosmetology. *Int J Res Dermatol.* 2020;6(2):288–91.
- Perez AG, Lana JF, Rodrigues AA, Luzo AC, Belangero WD, Santana MH, et al. Relevant aspects of centrifugation step in the preparation of Platelet-rich plasma. *ISRN Hematol.* 2014;p. 176060. doi:10.1155/2014/176060. eCollection 2014.
- Amable PR, Carias RBV, Teixeira MVT, Pacheco IC, Amaral R, Granjeiro JM, et al. Platelet-rich plasma preparation for regenerative medicine: Optimization and quantification of cytokines and growth factors. *Stem Cell Res Ther.* 2013;4(3):67. doi:10.1186/scrt218.
- Kahn RA, Cossette I, Friedman LI. Optimum centrifugation conditions for the preparation of platelet and plasma products. *Transfusion.* 1976;16(2):162–5.
- Slichter SJ, Harker LA. Preparation and storage of platelet concentrates. I. Factors influencing the harvest of viable platelets from whole blood. *Br J Haematol.* 1976;34(3):395–402.
- Landesberg R, Roy M, Glickman RS. Quantification of growth factor levels using a simplified method of platelet-rich plasma gel preparation. *J Oral Maxillofac Surg.* 2000;58(3):297–300.
- Jo CH, Roh YH, Kim JE, Shin S, Yoon KS. Optimizing platelet-rich plasma gel formation by varying time and gravitational forces during centrifugation. *J Oral Implantol.* 2013;39(5):525–32.
- Whitman DH, Berry RL, Green DM. Platelet gel: an autologous alternative to fibrin glue with applications in oral and maxillofacial surgery. *J Oral Maxillofac Surg.* 1997;55(11):1294–9.
- Gupta V, Parihar AS, Pathak M, Sharma VK. Comparison of Platelet-Rich Plasma Prepared Using Two Methods: Manual Double Spin

- Method versus a Commercially Available Automated Device. *Indian Dermatol Online J.* 2020;11(4):575–9.
14. Landesberg R, Moses M, Karpatkin M. Risks of using platelet rich plasma gel. *J Oral Maxillofac Surg.* 1998;56(9):1116–7.
  15. Landesberg R. Controlled Delivery of Growth Factors Derived From Platelet-Rich Plasma. *J Oral Maxillofac Surg.* 2006;64(9):87. doi:10.1016/j.joms.2006.06.159.
  16. Marx RE. Platelet-rich plasma: evidence to support its use. *J Oral Maxillofac Surg.* 2004;62(4):489–96.
  17. Haynesworth SE, Kadiyala S, Liang LN. Mitogenic stimulation of human mesenchymal stem cells by platelet release suggest a mechanism for enhancement of bone repair by platelet concentrates. In: Presented at the 48th Meeting of the Orthopedic Research Society, Boston, MA; 2002.
  18. Rughetti A, Giusti I, D'Ascenzo S, Leocata P, Carta G, Pavan A, et al. Platelet gel-released supernatant modulates the angiogenic capability of human endothelial cells. *Blood Transfus.* 2008;6(1):12–7.
  19. Ehrenfest DD, Rasmusson L, Albrektsson T. Classification of platelet concentrates: from pure platelet-rich plasma (P-PRP) to leucocyte- and platelet-rich fibrin (L-PRF). *Trends Biotechnol.* 2009;27(3):158–67.
  20. Oudelaar BW, Peerbooms JC, 't Veld R, Vochteloo A. Concentrations of blood components in commercial Platelet-rich plasma separation systems: A review of the literature. *Am J Sports Med.* 2019;47(2):479–87.
  21. Magalon J, Chateau AL, Bertrand B, Louis ML, Silvestre A, Giraudo L, et al. DEPA classification: a proposal for standardizing PRP use and a retrospective application of available devices. *BMJ Open Sport Exerc Med.* 2016;2(1):60. doi:10.1136/bmjsem-2015-000060.
  22. Braun HJ, Kim HJ, Chu CR, Dragoo JL. The effect of Platelet-rich plasma formulations and blood products on human synoviocytes: Implications for intra-articular injury and therapy. *Am J Sports Med.* 2014;42(5):1204–10.
  23. Uysal CA, Ertas NM. Platelet-Rich Plasma Increases Pigmentation. *J Craniofac Surg.* 2017;28(8):793. doi:10.1097/SCS.0000000000002893.
  24. Dhurat R, Sukesh MS. Principles and methods of preparation of Platelet-rich plasma: A review and author's perspective. *J Cutan Aesthet Surg.* 2014;7(4):189–97.
  25. Araki J, Jona M, Eto H, Aoi N, Kato H, Suga H, et al. Optimized preparation method of platelet-concentrated plasma and noncoagulating platelet-derived factor concentrates: Maximization of platelet concentration and removal of fibrinogen. *Tissue Eng Part C Methods.* 2012;18(3):176–85.
  26. Tamimi F, Montalvo S, Tresguerres I, Jerez LB. A comparative study of 2 methods for obtaining platelet-rich plasma. *J Oral Maxillofac Surg.* 2007;65(6):1084–93.
  27. Bausset O, Giraudo L, Veran J, Magalon J, Coudreuse JM, Magalon G, et al. Formulation and storage of Platelet-rich plasma homemade product. *Biores Open Access.* 2012;1(3):115–23.

## Author biography

**Zeal Jayeshbhai Rachchha**, Post Graduate Resident  
<https://orcid.org/0009-0003-3885-9558>

**Vaishali Sanket Makwana**, Associate Professor <https://orcid.org/0009-0002-4487-6832>

**Sandip Labhubhai Lathiya**, Ex-Post Graduate Student  
<https://orcid.org/0009-0003-9036-3074>

**Nehalben Manishbhai Pandya**, Professor and HOD  
<https://orcid.org/0009-0008-3822-9472>

**Cite this article:** Rachchha ZJ, Makwana VS, Lathiya SL, Pandya NM. A cross sectional study optimizing platelet rich plasma preparation using various centrifugation speeds and its effects on the platelet yield. *IP Indian J Clin Exp Dermatol* 2024;10(3):291-300.